

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07K 5/02, 7/02 C07D 307/33, A61K 37/64	A1	(11) International Publication Number: WO 94/03480 (43) International Publication Date: 17 February 1994 (17.02.94)
(21) International Application Number: PCT/US93/03589 (22) International Filing Date: 21 April 1993 (21.04.93) (30) Priority data: 07/922,889 31 July 1992 (31.07.92) US (60) Parent Application or Grant (63) Related by Continuation US 07/922,889 (CON) Filed on 31 July 1992 (31.07.92) (71) Applicant (for all designated States except US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US).		(72) Inventor; and (75) Inventor/Applicant (for US only): ROBINSON, Ralph, Pelton [US/US]; 30 Friar Tuck Drive, Gales Ferry, CT 06335 (US). (74) Agents: RICHARDSON, Peter, C. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US). (81) Designated States: CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: PEPTIDYL 4-AMINO-2,2-DIFLUORO-3-OXO-1,6-HEXANEDIOIC ACID DERIVATIVES AS ANTIINFLAMMATORY AGENTS <div style="text-align: center;"> <p style="margin-top: 10px;">(A)</p> </div> (57) Abstract <p>This invention relates to compounds of formula (A) which are derivatives of 4-amino-2,2-difluoro-3-oxo-1,6-hexanedioic acid and their pharmaceutically acceptable base salts. These derivatives and salts thereof are useful for inhibiting interleukin 1β converting enzyme and for treating inflammatory conditions in mammals. The invention also claims three (3) intermediates which are useful for the preparation of said derivatives and salts.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

PEPTIDYL 4-AMINO-2,2-DIFLUORO-3-OXO-1,6-HEXANEDIOIC
ACID DERIVATIVES AS ANTIINFLAMMATORY AGENTS

Background of the Invention

This invention is concerned with new antiinflammatory agents. In particular, this invention relates to compounds which are derivatives of 4-amino-2,2-difluoro-3-oxo-1,6-hexanedioic acid; to the pharmaceutically acceptable base salts of such derivatives; to methods of using such derivatives in inhibiting interleukin 1 β converting enzyme (ICE) and for treating inflammatory conditions in mammals, especially man; and to pharmaceutical compositions useful therefor.

Current therapies for arthritis are severely limited by the side effects of available drugs and their ineffectiveness beyond treatment for disease symptoms. The most widely used drugs are agents (the non-steroidal antiinflammatory drugs, NSAIDS) which inhibit the cyclooxygenase pathway of arachidonic acid metabolism. While these compounds are effective in controlling the symptoms of arthritis, they are not disease remittive. Furthermore, cyclooxygenase inhibition is invariably associated with the major side-effect of NSAID therapy, gastrointestinal irritation. Steroids are used in the more severe cases of arthritis and are very effective. However, long term therapy using steroids is seldomly tolerable. Second line antiinflammatory agents such as gold, penicillamine, chloroquine and methotrexate are also beset with side effect issues which severely limit their general utility.

Interleukin-1 (IL-1) has been strongly implicated as a key mediator of tissue damage in osteo- and rheumatoid arthritis. Lowering levels of IL-1 in a diseased joint would be expected to halt continued degeneration and perhaps allow joint repair to take place. One approach to reducing levels of IL-1 is to block the generation of mature IL-1 β from its biologically inactive precursor, pro-IL-1 β , by inhibition of the interleukin-1 β converting enzyme (ICE). This invention relates to a novel series of compounds which inhibit ICE. The compounds should act as disease remittive antiinflammatory agents and are not expected to elicit the side effects associated with NSAID therapy (due to cyclooxygenase inhibition), steroids or other treatments currently in use.

Peptidyl derivatives containing difluorostatone are described in: S. Thaisrivongs et al., J. Med. Chem., 1986, 29, 2080-2087 and K. Fearon et al., J. Med. Chem., 1987, 30, 1617-1622.


-2-

Summary of the Invention

This invention is concerned with new compounds which are useful for the treatment of diseases associated with elevated levels of interleukin-1 (IL-1). The compounds block the formation of biologically active mature IL-1 β from its precursor pro-IL-1 β by inhibiting interleukin 1 β converting enzyme (ICE).

The compounds of the present invention and their pharmaceutically acceptable salts are of the formula A

10




(A)

15

and the pharmaceutically acceptable base salts thereof

wherein A¹ is L-Pro-NR¹R² or -NR¹R², where R¹ and R² are independently selected from the group consisting of hydrogen, C₁-C₆ alkyl and benzyl; or R¹ and R² are taken together with the nitrogen to which they are attached and form

20



25

A² is selected from the group consisting of L-His, L-Cys, L-Cys(Me), L-Phe, L-Phe-R³, L-Val, L-Ala, L-Ile, L-Leu and L-Tyr;

A³ is selected from the group consisting of L-Val, L-Leu, L-Ile, L-Tyr, L-Phe and L-Phe-R³;

A⁴ is selected from the group consisting of a covalent bond, L-Phe, L-Phe-R³, L-Tyr, and L-Leu;

wherein R³ is attached to the aromatic ring of the phenylalanine and for each occurrence is selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ alkoxy, benzyl, fluoro, trifluoromethyl and chloro; and


35

-3-

Q¹ is selected from the group consisting of t-butoxycarbonyl, benzyloxycarbonyl, R⁴CO and phenylcarbonyl, wherein R⁴ is hydrogen, C₁-C₆ alkyl or benzyl.

5 The abbreviations used to denote the amino acids are well known and standard in the art and are as follows: Ala, alanine; Pro, proline; His, histidine; Cys, cystine; Cys (Me), methylcystine; Phe, phenylalanine; Val, valine; Ile, isoleucine; Leu, leucine; and Tyr, tyrosine.

A preferred group of compounds have the formula (A) wherein A¹ is
10



15

bond or L-Tyr, and Q¹ is t-butoxycarbonyl.

A more preferred group of compounds have the formula (A) wherein A¹, A³, A⁴ and Q¹ are the same as the preferred group of compounds and A² is L-Ala.

20 The invention provides a pharmaceutical composition comprising a compound of the formula (A), or a pharmaceutically acceptable base salt thereof, together with a pharmaceutically acceptable diluent or carrier.

The invention also provides a method of inhibiting interleukin 1 β converting enzyme (ICE) which comprises administering an effective amount of a compound of the
25 formula (A) or pharmaceutically acceptable base salt thereof, or a pharmaceutical composition as defined above.


The invention further provides a method of treating inflammatory conditions which comprises administering an effective amount of a compound of the formula (A) or pharmaceutically acceptable base salt thereof, or a pharmaceutical composition as
30 defined above. The term "inflammatory condition" includes arthritis, inflammatory bowel disease, psoriasis, allergic encephalitis, gingivitis, systemic lupus erythematosus, diabetes melitis, gout, septic shock and adult respiratory distress syndrome.

35

-4-


The present invention also claims the intermediate compounds, shown below, necessary for synthesizing the compounds of formula (A):

5




10

15



20

25



wherein A¹, A², A³, A⁴ and Q¹ are as previously defined.

30



Detailed Description of the Invention

The compounds of the present invention, having the formula (A) as defined above, are readily and generally prepared by the general method described below. Scheme I illustrates the reaction sequence that is followed to generate the compounds of the instant invention. The definitions of a¹, a², a³ and a⁴ are as follows: a¹ is L-Pro-NR¹R² or -NR¹R²; a² is N-benzyloxymethyl-L-His, S-benzyl-L-Cys, L-Cys(Me), L-Phe, L-

35

-5-

Phe-R³, L-Val, L-Ala, L-Ile, L-Leu or O-benzyl-L-Tyr; a³ is L-Val, L-Leu, L-Ile, O-benzyl-L-Tyr, L-Phe or L-Phe-R³; and a⁴ is L-Phe, L-Phe-R³, O-benzyl-L-Tyr or L-Leu. R¹, R², R³ and Q¹ are as defined hereinabove. The compounds of formula (I) and (II) have the following general structures:



20


25

30

35

-6-

Scheme 1



-7-

The starting material is the known enantiomerically pure aldehyde, (2S)-1-[(1,1-dimethylethyl)dimethylsilyl]-4-oxo-2-azetidinecarboxaldehyde (III) prepared according to the procedure of Labia and Morin (Chem. Lett. 1984, 1007). In the first step, III is reacted with $\text{BrZnCF}_2\text{CO}_2\text{Et}$ to give a mixture of the alcohol IV (ethyl [R-(R*,S*)]-1-[(1,1-dimethylethyl)dimethylsilyl]- α,α -difluoro- β -hydroxy-4-oxo-2-azetidinepropanoate) and its C β epimer (ethyl [S-(S*,S*)]-1-[(1,1-dimethylethyl)dimethylsilyl]- α,α -difluoro- β -hydroxy-4-oxo-2-azetidinepropanoate). The reaction is typically carried out in tetrahydrofuran as solvent although other ether solvents such as diethyl ether and 1,2-dimethoxyethane can be used. The $\text{BrZnCF}_2\text{CO}_2\text{Et}$ is normally formed in situ by reaction of ethyl bromodifluoroacetate with zinc powder (activated by washing with dilute mineral acid and drying). The formation of $\text{BrZnCF}_2\text{CO}_2\text{Et}$ can be accelerated greatly by ultrasound irradiation either by carrying out the reaction in an ultrasound bath or by the use of an immersed ultrasound probe. Thus the reaction is normally carried out by dissolving III (about 1 equivalent) and ethyl bromodifluoroacetate (about 1.1 to 10 equivalents, typically 2 equivalents) in the reaction solvent, adding Zn powder (about 2 to 100 equivalents, typically 3 equivalents) and immersing the reaction vessel in an ultrasound bath. The reaction temperature using ultrasound can vary between about 0°C and the boiling point of the solvent (provided this is less than 100°C). A reaction temperature of approximately 35°C is normally used with ultrasound irradiation; under these conditions the reaction time is about 1 hour although shorter (10 minutes) or longer reaction times (12 hours) can be used. The reaction may alternatively be carried out without ultrasonic irradiation. Under these conditions, higher reaction temperatures may be required although the upper limit for temperature still applies (100°C). Another means of carrying out the reaction is to pre-form the $\text{BrZnCF}_2\text{CO}_2\text{Et}$ by the reaction of zinc powder (about 3 equivalents) with ethyl bromodifluoroacetate (about 3 equivalents) and iodine (about 0.05 equivalents) in an ether solvent (for example tetrahydrofuran, diethyl ether, or 1,2-dimethoxyethane) using heat or ultrasound (Altenburger and Schirlin, Tetrahedron Letters 1991, 32, 7255). After formation of the organozinc reagent, a solution of III, dissolved in the same solvent being used for the reaction, is added dropwise and when addition is complete, ultrasonication or heating is continued for up to about 12 hours. Yet another procedure is to add a solution of III and ethyl bromodifluoroacetate (about 10 equivalents) in an ether solvent (for example tetrahydrofuran, diethyl ether, or 1,2-dimethoxyethane) to the preformed low-valent

-8-

titanium species formed by the reaction of zinc powder (about 1 equivalent) with titanium tetrachloride (about 0.6 equivalents) (Parris et al, Biochemistry, 1992, in press) 5 in the same solvent. Regardless of the manner in which the reaction is carried out, the workup is the same. The reaction is poured into water or a saturated aqueous solution of ammonium chloride and extracted with an immiscible organic solvent preferably diethyl ether or ethyl acetate. After drying, evaporation of the solvent leaves the crude mixture of products. The desired product **IV** is isolated by flash chromatography on 10 silica gel eluting with 3:7 ethyl acetate/hexane; the more polar C β epimer is eluted after **IV**.

The next step involves the removal of the t-butyldimethylsilyl (TBDMS) protecting group from the β -lactam nitrogen. This reaction is normally carried out by treatment of a solution of **IV** in acetonitrile with an excess of HF/pyridine complex. Reaction 15 temperatures between -20°C and 50°C are acceptable, with about 0°C being preferred. At this temperature the reaction time is about 1.5 hours although reaction times may vary between 10 minutes and 12 hours. The reaction is typically worked up by dilution with water and repeated extraction with ethyl acetate. After drying and removal of solvent, the crude product **V** remains and is typically of sufficient purity for use in the 20 next step. Other procedures well precededented in the literature may also be applicable to carrying out this reaction, for example HCl in methanol/water (Salzmann et al. J. Am. Chem. Soc., 1980, 102, 6161).

The ethyl ester **V** is subsequently converted to the corresponding amide **VI**. For the formation of amides derived from primary and secondary amines ($a^1 = NR^1R^2$), the 25 conversion is carried out by direct reaction between **V** (about 1 equivalent) and the amine HNR^1R^2 (1 to 20 equivalents, typically 2 equivalents). The preferred solvent for the reaction is methylene chloride although many other solvents including toluene, chloroform, ethanol, N,N-dimethylformamide, and ethyl acetate may be used. Reaction temperatures vary between about 0°C and 80°C; at higher temperatures unwanted side- 30 reactions involving attack by the amine on the β -lactam may occur. The usual reaction temperature in methylene chloride is about 20°C; and the reaction time is typically about 1.5 hours, although the time can be varied from 1 hour to 1 day. Workup of the reaction is normally carried out by evaporating the solvent followed by flash chromatography on silica gel or diluting with ethyl acetate, washing with dilute mineral 35 acid and evaporating the solvent. For the formation of L-proline amides **VI**

(a¹ = L-ProNR¹R²), amide bond formation is carried out by prior hydrolysis of the ethyl ester group to give the corresponding carboxylic acid followed by coupling with the appropriate L-proline derivative (H-L-ProNR¹R²). The hydrolysis step is carried out by the action of MOH (1 to 1.5 equivalents, typically about 1.1 equivalents; M=Na, K, or Li) in a 1:10 mixture of H₂O and tetrahydrofuran at about 0°C followed by neutralizing with aqueous mineral acid, extracting with ethyl acetate, and removal of solvent. Numerous methods exist which can be used in the subsequent amide bond forming step; these are well detailed in the literature and include coupling with bis(2-oxo-3-oxazolidinyl)phosphinic chloride (Rich et al., J. Org. Chem., 1990, 55, 2895), diethylphosphoryl cyanide (Yamada et al., Tetrahedron Lett., 1973, 14, 1595), N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline (Belleau and Malek, J. Am. Chem. Soc., 1968, 90, 1651), dicyclohexylcarbodiimide (Konig and Geiger Ber., 1970, 103, 788) and mixed anhydride formation (Vaughn and Osato J. Am. Chem. Soc., 1951, 73, 5553).

The next step involves coupling of the intermediate **VI** (about 1 equivalent) with an N-t-butoxycarbonyl α -amino acid N-hydroxysuccinimide ester (1 to 10 equivalents, typically about 1.1 equivalents) with accompanying β -lactam ring opening and lactonization to give **VII**. The reaction is carried out in the presence of a tertiary amine base (1 to 10 equivalents, typically about 2 equivalents) such as triethylamine or diisopropylethylamine. An inert solvent such as chloroform, toluene, ethyl acetate or methylene chloride (preferably methylene chloride) is used. Reaction temperatures from 0°C to 80°C can be used depending on the solvent. The normal temperature used in methylene chloride is about 20°C, and the reaction time is typically about 18 hours, although the time can be varied from 12 hours to 3 days. The product **VII** is normally isolated by evaporating the solvent followed by flash chromatography on silica gel.

Peptide chain extension reactions analogous to the formation of lactone **VIII** from **VII** are described in the literature and are well known to those skilled in the art of peptide synthesis. The N-t-butoxycarbonyl protecting group is first removed by treating a solution of **VII** in methylene chloride with an excess of trifluoroacetic acid (TFA) at about 0°C. The volatile solvent and TFA are removed in vacuo and the intermediate TFA amine salt is coupled to an amino acid derivative of structure BOC-a³-OH, BOC-a³-OX, Z-a³-OH or Z-a³-OX (where BOC is t-butoxycarbonyl, Z is benzyloxycarbonyl, X is N-succinimidyl or pentafluorophenyl) in the presence of a tertiary amine base (e.g. triethylamine or diisopropylethylamine) to give **VIII** (Q¹ is t-butoxycarbonyl or

-10-

benzyloxycarbonyl). The preferred solvent for the coupling step is methylene chloride, although other solvents, e.g. chloroform or N,N-dimethylformamide can be used.

5 For the preparation of compounds of formula I, (Q¹ is t-butoxycarbonyl or benzyloxycarbonyl), an intermediate **VIII** (Q¹ is t-butoxycarbonyl or benzyloxycarbonyl) is next converted to a benzyl ester **IX** (Q¹ is t-butoxycarbonyl or benzyloxycarbonyl). This is accomplished by hydrolyzing the lactone function by treating **VIII** (Q¹ is t-butoxycarbonyl or benzyloxycarbonyl) with MOH (1 to 1.5 equivalents, preferably about
10 1.1 equivalents; M is Na, K, or Li, preferably Li) in a 1:5 mixture of H₂O and tetrahydrofuran at 20°C. The time for this reaction is typically about 2 hours. The solvents are evaporated to leave the metal salt of the corresponding γ-hydroxyester as a solid which is dried under vacuum. The salt is dissolved in dry N,N-dimethylformamide and the resulting solution is treated with benzyl bromide (1 to 10
15 equivalents, preferably about 1.5 equivalents). The solution is stirred for 1 to 12 hours (typically 5 hours) at about 20°C and is then poured into water. Other reaction temperatures between 0°C and 80°C can be used. The mixture is extracted with ethyl acetate and the combined ethyl acetate fractions are dried and concentrated to leave the crude product mixture. The intermediate **IX** (Q¹ is t-butoxycarbonyl or
20 benzyloxycarbonyl) is then purified by flash chromatography of the mixture on silica gel, typically using ethyl acetate or some combination of ethyl acetate and hexane as eluant.

For the preparation of compounds of formula I, (Q¹ is R¹CO or PhCO), an intermediate **VIII** (Q¹ is t-butoxycarbonyl) is treated with trifluoroacetic acid (TFA) as in
25 the conversion of **VII** to **VIII**. The resulting TFA salt of the corresponding amino compound is then treated with an acid chloride of the formula R¹COCl or PhCOCl or an anhydride of the formula (R¹CO)₂O or (PhCO)₂O in the presence of a tertiary amine such as triethylamine to give **VIII** (Q¹ is R¹CO or PhCO). This is subsequently transformed into **IX** (Q¹ is R¹CO or PhCO) by the same procedure as for the conversion
30 of **VIII** (Q¹ is t-butoxycarbonyl or benzyloxycarbonyl) to **IX** (Q¹ is t-butoxycarbonyl or benzyloxycarbonyl).

The final reaction sequence to provide compounds of formula I is as follows. The alcohol function of an intermediate **IX** is first oxidized to the ketone using 1,1,1-triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1H)-one according to the procedure described
35 by Linderman (Tetrahedron Lett., 1987, 28, 4259). Next, the benzyl ester function is

-11-

cleaved to give I by catalytic hydrogenolysis using palladium on charcoal. The preferred solvent for the hydrogenolysis is ethanol although other solvents such as ethyl acetate or acetic acid may be used. The preferred pressure of hydrogen is about 3 atmospheres although the range may vary between 1 and 50 atmospheres. The reaction temperature is typically about 20°C. The product I is isolated by filtration to remove the catalyst followed by evaporation of solvent. For the preparation of compounds of formula I where Q¹ is benzyloxycarbonyl, it is often preferable to hydrolyze the benzyl ester by treatment of IX (Q¹ is benzyloxycarbonyl) with MOH (1 to 1.5 equivalents, preferably about 1.1 equivalents; M = Na, K, or Li, preferably Li) in a 1:5 mixture of H₂O and tetrahydrofuran at about 20°C. The compound of formula I (Q¹ is benzyloxycarbonyl) is then isolated by neutralization of the reaction mixture with mineral acid, extraction with ethyl acetate, removal of solvent and chromatography on silica gel.

To synthesize compounds of formula II, an intermediate VIII (Q¹ is t-butoxycarbonyl) is extended to an intermediate X (Q¹ is t-butoxycarbonyl or benzyloxycarbonyl) by the same procedure used in the conversion of VII to VIII (Q¹ is t-butoxycarbonyl or benzyloxycarbonyl) using BOC-a⁴-OH, BOC-a⁴-OX, Z-a⁴-OH or Z-a⁴-OX (where BOC is t-butoxycarbonyl, Z is benzyloxycarbonyl, X is N-succinimidyl or pentafluorophenyl). For the synthesis of compounds of formula II (Q¹ is R¹CO or PhCO), an intermediate X (Q¹ is t-butoxycarbonyl) is treated with trifluoroacetic acid (TFA). The resulting TFA salt of the corresponding amino compound is then treated with an acid chloride of the formula R¹COCl or PhCOCl or an anhydride of the formula (R¹CO)₂O or (PhCO)₂O in the presence of a tertiary amine such as triethylamine to give X (Q¹ is R¹CO or PhCO). The final conversions of X to XI and then to II are carried out as described for the sequence VIII to IX to I. For the preparation of compounds of formula II where Q¹ is benzyloxycarbonyl, it is often preferable to cleave the benzyl ester by treatment of XI with MOH as for the synthesis of compounds of formula I (Q¹ is benzyloxycarbonyl).

Compounds where a² is L-His or L-Cys, L-Tyr, and/or a⁴ is L-Tyr require side chain protection. These amino acids are introduced with the following side chain protecting groups: L-His, N-benzyloxymethyl; L-Cys, S-benzyl; L-Tyr, O-benzyl. With L-His the side chain protecting group is removed during the final hydrogenolysis of the

-12-

benzyl ester giving I or II. If Q¹ is benzyloxycarbonyl, loss of this group may occur necessitating treatment of the product with benzyl chloroformate to re-introduce the
5 benzyloxycarbonyl group. If L-Cys and/or L-Tyr are introduced, the benzyl side chain protecting groups of these amino acids are removed from the intermediates IX or XI by treatment with neat HF. These conditions also serve to cleave the benzyl ester, the N-benzyloxymethyl on L-His (if present), and/or the benzyloxycarbonyl group at Q¹ (if present) or the t-butoxycarbonyl group at Q¹ (if present). If the Q¹ group is lost, i.e.
10 when Q¹ is t-butoxycarbonyl or benzyloxycarbonyl, the group may be re-introduced by treatment with di-t-butylidicarbonate or benzyl chloroformate respectively.

The ability of the compounds of this invention to inhibit interleukin 1 β converting enzyme (ICE) and, consequently, demonstrate the compounds' effectiveness for treating inflammatory diseases is shown by the following in vitro assay. Other procedures for
15 purification and assaying (ICE) are shown in Black et al., Febs Letters, 247 2, pp. 386-390, 1989, and Thornberry et al., Nature, 356, pp. 768-774, 1992.

Cell culture and lysates. Human monocyte cell line, THP-1 (ATCC-TIB 202) is grown in RPMI media 1640 (Gibco BRL Gaithersburg, MD 20877) with 10% fetal bovine serum, harvested by centrifugation, washed twice in Dulbecco's PBS dithiothreitol
20 without Ca⁺⁺ ethylene diamine tetraacetic acid and Mg⁺⁺, and resuspended in buffer (10 mM Tris-HCl pH 7.8, 5 mM DTT (dithiothreitol), 1 mM EDTA, 1 mM PMSF (phenylmethyl sulfonyl fluoride), 1 μ g/ml pepstatin, and 1 μ g/ml leupeptin) at 1-3 x 10⁶ cells per ml. Cells are frozen at -70°C until use and then lysed by thawing. Lysates are cleared by centrifugation at 20,000 x g for 1 hour followed by 120,000 x g for 1 hour.

25 Partial purification of ICE activity by ion-exchange chromatography. ICE activity is purified from THP-1 cell lysates by three steps of ion exchange chromatography.

(A) THP-1 cell lysate (1.5 L) is chromatographed by ion-exchange on Q-Sepharose Fast Flow (Pharmacia LKB Biotechnology Piscataway, NJ 08854) in buffer A (20 mM Tris pH 7.8, 5 mM EDTA, 1 mM PMSF, 1 μ g/ml pepstatin, and 1 μ g/ml leupeptin). ICE activity
30 is eluted with a gradient of NaCl in buffer A. (B) The active pool from A is chromatographed by anion exchange on MonoQ monobeads (Pharmacia LKB Biotechnology Piscataway, NJ 08854) in buffer A and activity eluted in a NaCl gradient. (C) The active pool from B is chromatographed by cation exchange on monoS monobeads (Pharmacia LKB Biotechnology Piscataway, NJ 08854) in buffer B (25 mM
35 NaMES (2-[N-morpholino]ethanesulfonic acid, hemisodium salt) pH 6.9, 5mM DTT, 1

-13-

mM EDTA, 10% glycerol, 1 mM PMSF, 1 μ g/ml pepstatin, and 1 μ g/ml leupeptin).

Activity is eluted in a NaCl gradient. The active pool from C is used in the subsequent
5 assay to measure the ICE inhibitory activity of the compounds comprising this invention.

ICE Assay. ICE activity is assayed by combining 10 μ l partially purified enzyme
fraction containing 10 mM Tris-HCl (pH 7.8) with protease inhibitors (1 mM PMSF, 1
 μ g/ml pepstatin, and 1 μ g/ml leupeptin), with test compound in a 40 μ l volume (yielding
a 20% DMSO concentration after substrate is added). The reaction is initiated by the
10 introduction of [35 S]-metabolically labelled human peripheral blood monocyte proteins
(containing [35 S] labelled pro-IL-1 β substrate) in a total volume of 50 μ l and terminated
10 minutes later by the addition of 10 μ l of 0.9 M NaCl. The cleaved and uncleaved
forms of IL-1 β are immunocaptured by the addition of 5 μ l of polyclonal anti-IL-1 β
antibody (Cistron Biotechnology, Box 2004, 10 Bloomfield Avenue, Pine Brook, NJ
15 07058, product #02-1100) in 50 μ l PBS (Dulbecco's Phosphate Buffered Saline) (pH 8)
plus 0.1% SDS (Sodium Dodecyl Sulfate Detergent), 0.1% Triton X-100 (Nonionic
detergent Sigma Chemical Co., P.O. Box 14508, St. Louis, MO 63178), 0.1% Nonidet
P40 (NP-40, Nonionic Detergent Sigma Chemical Co., P.O. Box 14508, St. Louis, MO
63178) and further incubation for 4 hours at 4°C. Immune complexes are insolubilized
20 by the addition of 25 μ l of a 50% slurry of Protein-A Sepharose, and incubation
overnight at 4°C. The insolubilized complexes are recovered by centrifugation
(Eppendorf microfuge), and washed three times using a 10X volume of PBS (pH 8) with
0.1% Triton X-100, 0.1% SDS, 0.1% NP-40. After the last wash, a volume of 25 μ l
containing 2% SDS, 200 mM dithiothreitol in 0.125 M Tris buffer (pH 6.8) (2X Laemmli
25 sample buffer [0.125M Tris-HCl pH 6.8, 4% SDS, 20% Glycerol, 10% 2-mercaptoethanol,
0.002% Bromophenol Blue]) with 20% glycerol are added to the pellet. The immune
complexes are released from the Protein A and the [IL1-anti-IL1] complexes dissociated
by immersing the tubes in boiling water. Two minutes later, the tubes are chilled in ice,
and centrifuged to pellet the protein A Sepharose.

30 After clarification by centrifugation, a 25 μ l aliquot is applied to a 10-20%
gradient of Laemmli SDS PAGE (Integrated Separation Systems, One Westinghouse
Plaza, Hyde Park, MA 02136) and electrophoresed at 50 mA per gel for approximately
75 minutes. After removal, the gels are fixed in 10% HOAc:30% MeOH for 60 minutes
at ambient, and bathed in AMPLIFY (Amersham Co., Arlington Heights, IL 60005) for 30
35 minutes at ambient. The gels are then dried on a vacuum drier (Hoeffer) for 90 minutes

-14-

at 60°C, and developed using Kodak XAR film (Parker X-Ray, 260 Governor Street, E. Hartford, CT 06108) at -70°C.

5 Bands corresponding to pro- and mature forms of IL-1 β are quantitated by elution using 0.5 ml of 0.1 M Tris (pH 8), 20 mM CaCl₂, and 10 mg/ml pronase incubated at 56° for 4 hours. After the addition of 4 ml of Ready-Safe (Liquid Scintillation Cocktail) (Beckman Instruments Inc., Fullerton, CA 92034), the digests are counted using a liquid scintillation counter.

10 The pharmaceutically acceptable base salts of the compounds of the formula (A) are those formed from bases which form non-toxic base salts, for example the sodium, potassium and ammonium salts.

For administration to humans in the curative or prophylactic treatment of inflammatory conditions using a compound of formula (A) above, oral dosages of the
15 compounds are generally in the range from 2-100 mg daily for an average adult patient (70 kg). Thus for a typical adult patient, individual tablets or capsules containing from 1 to 10 mg of active compound, in a suitable pharmaceutically acceptable vehicle or carrier, are used. Dosages for intravenous administration are typically within the range 1 to 10 mg per single dose as required. In practice the physician will determine the
20 actual dosage which will be most suitable for an individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are exemplary of the average case but there can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

25 For human use, the compounds of the formula (A) can be administered alone, but will generally be administered in an admixture with a pharmaceutical carrier selected with regard to the intended route of administration and standard pharmaceutical practice. For example, they may be administered orally in the form of tablets containing such excipients as starch or lactose, or in capsules or ovals either alone or in
30 admixture with excipients, or in the form of elixirs or suspensions containing flavoring or coloring agents. They may be injected parenterally, for example, intravenously, intramuscularly or subcutaneously. For parenteral administration, they are best used in the form of a sterile aqueous solution which may contain other substances, for example, enough salts or glucose to make the solution isotonic.

35

-15-

The present invention is illustrated by the following examples, but it is not limited to the details thereof.

5

EXAMPLE 1

[S]-N-[(1,1-Dimethylethoxy)carbonyl-L-Valyl-N-[3,3-Difluoro-2,4-Dioxo-1-Carboxymethyl-4-(1-Pyrrolidiny)Butyl]-L-Alaninimide

- A. Ethyl [S-(S*,S*)]-1-[(1,1-dimethylethyl)dimethylsilyl]- α,α -difluoro- β -hydroxy-4-oxo-2-azetidinepropanoate and ethyl [R-(R*,S*)]-1-[(1,1-dimethylethyl)dimethylsilyl]- α,α -difluoro- β -hydroxy-4-oxo-2-azetidinepropanoate

To a solution of freshly prepared crude (2S)-1-[(1,1-dimethylethyl)dimethylsilyl]-4-oxo-2-azetidinecarboxaldehyde (10.3 g, 48.3 mmol) and ethyl bromodifluoroacetate (25.0 g, 123 mmol) in THF (100 ml) was added Zn powder (10.5 g, 161 mmol). The reaction flask was placed in a sonicating bath at 35°C for 40 minutes with occasional manual agitation. The mixture was then poured into ice/H₂O and the resulting slurry was filtered through celite, washing well with ether. The aqueous layer was separated and extracted with ether. The mixture was chromatographed on silica gel using 3:7 ethyl acetate/hexane as eluant. Complete separation of the epimeric products was not achieved. Fractions containing only the less polar product provided a solid which was triturated with hexane leaving pure ethyl [R-(R*,S*)]-1-[(1,1-dimethylethyl)dimethylsilyl]- α,α -difluoro- β -hydroxy-4-oxo-2-azetidinepropanoate as white crystals (5.17 g, 31.5%), m.p. 91-93°C, $[\alpha]_D^{20}$ -10.6° (c 1.13, CHCl₃), ν_{\max} 3578, 1747 cm⁻¹. MS (FAB): m/z 338. Elemental Anal. Calc'd for C₁₄H₂₅F₂NO₄Si: C, 49.83; H, 7.47; N, 4.12. Found: C, 49.90; H, 7.28; N, 4.15.

Fractions containing only the more polar product provided a solid which was triturated with hexane leaving pure ethyl [S-(S*,S*)]-1-[(1,1-dimethylethyl)dimethylsilyl]- α,α -difluoro- β -hydroxy-4-oxo-2-azetidinepropanoate as white crystals (1.26 g, 7.7%), mp 101-103°C, $[\alpha]_D^{20}$ -54.8° (c 1.70, CHCl₃), ν_{\max} 3669, 1738 cm⁻¹. MS (FAB): m/z 338. Elemental Anal. Calc'd for C₁₄H₂₅F₂NO₄Si: C, 49.83; H, 7.47; N, 4.12. Found: C, 49.93; H, 7.39; N, 4.15.

- B. [R-(R*,S*)]- α,α -Difluoro- β -hydroxy-4-oxo-2-azetidinepropanoate

To a solution of [R-(R*,S*)]-1-[(1,1-dimethylethyl)dimethylsilyl]- α,α -difluoro- β -hydroxy-4-oxo-2-azetidinepropanoate (1.0 g, 2.96 mmol) in CH₃CN (20 ml) at 0°C was added HF/pyridine (1 ml). After stirring for 40 minutes at 0°C, more HF/pyridine (1 ml)

-16-

was added; stirring was continued for an additional 30 minutes. The mixture was diluted with H₂O and then extracted with ethyl acetate (3 x). The combined ethyl acetate extracts were dried and concentrated to afford [R-(R*,S*)]- α,α -difluoro- β -hydroxy-4-oxo-2-azetidinepropanoate as a pale orange oil (667 mg, 100%). ¹HNMR: 1.32 (t, 3 H, J=7.1 Hz, CH₂Me), 2.90 (br d, 1 H, J=15.0 Hz, CHCON), 3.07 (ddd, J=1.5, 4.9, 15.0 Hz, CHCON), 3.98 (e, 1 H, CHCH₂), 4.09 (dt, 1 H, J=5.0, 18 Hz, CHCF₂), 4.33 (q, 2 H, J=7.1 Hz, CH₂Me), 4.70 (br s, 1 H, OH), 6.99 (br s, 1 H, NH).

10 C. [R-(R*,S*)]-1-[3-(4-Oxo-2-azetidiny)-2,2-difluoro-3-hydroxy-1-oxopropyl]pyrrolidine

[R-(R*,S*)]- α,α -Difluoro- β -hydroxy-4-oxo-2-azetidinepropanoate was dissolved in CH₂Cl₂ (5 ml). The solution was cooled to 0°C and pyrrolidine (0.5 ml, 6.0 mmol) was added. m.p. 104-106°C, [α]_D²⁰ +10.0° (c 2.08, CH₂Cl₂), ν_{\max} 3416, 1766, 1646
15 cm⁻¹. MS (EI): m/z 249 (M + H⁺). Elemental Anal. Calc'd for C₁₀H₁₄F₂N₂O₃: C, 48.39; H, 5.68; N, 11.29. Found: C, 48.51; H, 5.68; N, 11.24.

D. (2R-cis)-N-[(1,1-Dimethylethoxy)carbonyl]-N-[2-[1,1-difluoro-2-oxo-2-(1-pyrrolidinyl)ethyl]tetrahydro-5-oxo-3-furanyl]-L-alaninimide

To a solution of [R-(R*,S*)]-1-[3-(4-oxo-2-azetidiny)-2,2-difluoro-3-hydroxy-1-oxopropyl]-pyrrolidine (158 mg, 0.637 mmol) and triethylamine (0.18 ml, 1.29 mmol) in CH₂Cl₂ (4 ml) at 25°C was added N-(tert-butoxycarbonyl)-L-alanine N-hydroxy-succinimide ester (200 mg, 0.698 mmol). The mixture was stirred at 25°C for 18 hours and then concentrated to leave an oil. This was chromatographed on silica gel eluting with 3:1 ethyl acetate/hexane as eluant. Fractions containing only (2R-cis)-N-[(1,1-
25 dimethylethoxy)carbonyl]-N-[2-[1,1-difluoro-2-oxo-2-(1-pyrrolidinyl)ethyl]tetrahydro-5-oxo-3-furanyl]-L-alaninimide yielded an oil (175 mg, 66%), [α]_D²⁰ -45.9° (c 3.50, CH₂Cl₂), ν_{\max} 3428, 3324, 1802, 1693, 1650 cm⁻¹. MS (FAB): m/z 420.

E. (2R-cis)-N-[(1,1-Dimethylethoxy)carbonyl]-L-valyl-N-[2-[1,1-difluoro-2-oxo-2-(1-pyrrolidinyl)ethyl]tetrahydro-5-oxo-3-furanyl]-L-alaninimide

30 To a solution of (2R-cis)-N-[(1,1-dimethylethoxy)carbonyl]-N-[2-[1,1-difluoro-2-oxo-2-(1-pyrrolidinyl)ethyl]tetrahydro-5-oxo-3-furanyl]-L-alaninimide (260 mg, 0.620 mmol) in CH₂Cl₂ (4 ml) cooled in an ice bath was slowly added TFA (4 ml). The resulting mixture was stirred for 2 hours at 0°C at which point the solvents were evaporated to leave an oil. This was dissolved in CH₂Cl₂ and the solution was cooled
35 in an ice bath. Diisopropylethylamine (0.20 ml, 1.15 mmol) and BOC-L-valine-N-

-17-

hydroxysuccinimide ester (290 mg, 0.922 mmol) were added sequentially. After stirring for 4 hours at 25°C, the mixture was diluted with ethyl acetate and washed with 1N HCl.

5 The aqueous wash was extracted with ethyl acetate. The combined ethyl acetate fractions were washed with saturated NaHCO₃, dried and concentrated to leave an oil which was chromatographed on silica gel eluting with 4:1 ethyl acetate/hexane. Fractions containing only (2R-cis)-N-[(1,1-dimethylethoxy)carbonyl]-L-valyl-N-[2-[1,1-difluoro-2-oxo-2-(1-pyrrolidinyl)ethyl]tetrahydro-5-oxo-3-furanyl]-L-alaninimide yielded an
10 oil, (231 mg, 72%), $[\alpha]_D^{20}$ -47.8° (c 1.38, CHCl₃), ν_{\max} 3416, 3326, 1800, 1702 (sh), 1695, 1656 cm⁻¹. High Resolution MS Calc'd for C₂₇H₃₇F₂N₄O₇: 519.2630. Found: 519.2620.

F. [R-(R*,S*)]-N-[(1,1-Dimethylethoxy)carbonyl]-L-valyl-N-[3,3-difluoro-2-hydroxy-4-oxo-1-[2-oxo-2-(phenylmethoxy)ethyl]-4-(1-pyrrolidinyl)butyl]-L-alaninimide

15 A solution of (2R-cis)-N-[(1,1-dimethylethoxy)carbonyl]-L-valyl-N-[2-[1,1-difluoro-2-oxo-2-(1-pyrrolidinyl)ethyl]tetrahydro-5-oxo-3-furanyl]-L-alaninimide in MeOH (1.5 ml) and THF (1.5 ml) was cooled in an ice bath at 0°C and aqueous 1.31 M LiOH solution (0.25 ml, 0.33 mmol) was added. The mixture was stirred at 0°C for 1 hour and then at 20°C for 2 hours until disappearance of starting material was complete as determined by TLC
20 (silica gel, ethyl acetate eluant). After evaporation of the solvents, the residue was dried under vacuum for 18 hours to afford a white solid. This was dissolved in dry DMF (3 ml) and benzyl bromide (0.055 ml, 0.46 mmol) was added. After stirring at room temperature for 4.5 hours, the solution was poured into water and the resulting mixture was extracted with ethyl acetate (3 x). The combined ethyl acetate extracts gave a pale
25 yellow oil on drying and evaporation. This was chromatographed on silica gel eluting with 4:1 ethyl acetate/hexane to afford [R-(R*,S*)]-N-[1,1-dimethylethoxy)carbonyl]-L-valyl-N-[3,3-difluoro-2-hydroxy-4-oxo-1-[2-oxo-2-(phenylmethoxy)ethyl]-4-(1-pyrrolidinyl)butyl]-L-alaninimide as a clear oil, (135 mg, 71%), $[\alpha]_D^{20}$ -23.8° (c 1.14, CHCl₃), ν_{\max} 3423, 1728 (sh), 1708 (sh), 1649 cm⁻¹. HRMS Calc'd for C₃₀H₄₅F₂N₄O₈:
30 627.3208. Found: 627.3151.

G. [S]-N-[(1,1-Dimethylethoxy)carbonyl]-L-valyl-N-[3,3-difluoro-2,4-dioxo-1-[2-oxo-2-(phenylmethoxy)ethyl]-4-(1-pyrrolidinyl)butyl]-L-alaninimide

To a solution of [R-(R*,S*)]-N-[(1,1-dimethylethoxy)carbonyl]-L-valyl-N-[3,3-difluoro-2-hydroxy-4-oxo-1-[2-oxo-2-(phenylmethoxy)ethyl]-4-(1-pyrrolidinyl)butyl]-L-
35 alaninimide (129 mg, 0.206 mmol) in CH₂Cl₂ (5 ml) at 20°C was added the Dess-Martin

-18-

reagent (1,1,1-triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1H)-one) (320 mg, 0.75 mmol). The resulting heterogeneous mixture was stirred at 20°C for 5 hours and was then
 5 quenched by addition of ethyl acetate and a solution of sodium thiosulfate (1.2 g) in saturated NaHCO₃ solution (10 ml). After stirring for about 15 minutes when all solids had dissolved, the aqueous layer was separated and extracted with ethyl acetate. The combined ethyl acetate layers were dried and evaporated to leave [S]-N-[(1,1-dimethylethoxy)carbonyl]-L-valyl-N-[3,3-difluoro-2,4-dioxo-1-[2-oxo-2-
 10 (phenylmethoxy)ethyl]-4-(1-pyrrolidinyl)butyl]-L-alaninimide as a clear oil (126 mg, 98%). Having a high degree of purity by ¹H NMR, the material was used immediately in the next step. ¹H NMR: 0.87 (d, 3 H, J=6.8 Hz, CHMe), 0.93 (d, 3 H, J=6.8 Hz, NCHMe), 1.33 (d, 3 H, J=7.0 Hz, NCHMe), 1.42 (s, 9 H, t-Bu), 1.81-2.00 (m, 4 H, 2 x CH₂CH₂N), 2.10 (m, 1 H, CHMe₂), 2.97 (ABX m, 2 H, CH₂CO₂), 3.47 (e, 2 H, 2 x CH₂CHN), 3.62 (m,
 15 2 H, 2 x CH₂CHN), 3.95 (m, 1 H, NCHi-Pr), 4.52 (m, 1 H, NCHMe), 5.09 (AB d, 2 x 1 H, J=12.2 Hz, PhCH₂), 5.15-5.28 (m, 2 H, CHCH₂, BOCNH), 6.82 (br d, 1 H, J=7.4 Hz, NH), 7.30-7.41 (m, 6 H, Ph, NH). HRMS (FAB) Calc'd for C₃₀H₄₃F₂N₄O₈: 625.3051. Found: 627.3068.

H. [S]-N-[(1,1-Dimethylethoxy)carbonyl]-L-valyl-N-[3,3-difluoro-2,4-dioxo-1-carboxymethyl-4-(1-pyrrolidinyl)butyl]-L-alaninimide
 20


To a solution of [S]-N-[(1,1-dimethylethoxy)carbonyl]-L-valyl-N-[3,3-difluoro-2,4-dioxo-1-[2-oxo-2-(phenylmethoxy)ethyl]-4-(1-pyrrolidinyl)butyl]-L-alaninimide (124 mg, 0.198 mmol) in EtOH (25 ml) was added 10% Pd on charcoal. The mixture was hydrogenated at 3 atmospheres pressure for 5 hours using a Parr shaker. After
 25 removal of the catalyst by filtration through celite, the solvent was evaporated. The residue was chromatographed on silica gel using 1:5:54 AcOH/MeOH/CHCl₃ as eluant affording [S]-N-[(1,1-dimethylethoxy)carbonyl]-L-valyl-N-[3,3-difluoro-2,4-dioxo-1-carboxymethyl-4-(1-pyrrolidinyl)butyl]-L-alaninimide as a clear oil (103 mg, 97%). ν_{\max} 3422, 1812, 1684, 1649 cm⁻¹. HRMS (FAB) Calc'd for C₂₃H₃₇F₂N₄O₈ (MH⁺) = 535.2581. Found: 535.2597.
 30

-19-

CLAIMS

1. A compound of the formula

5




(A)

10

and the pharmaceutically acceptable base salts thereof

wherein A¹ is L-Pro-NR¹R² or -NR¹R², where R¹ and R² are independently selected from the group consisting of hydrogen, C₁-C₆ alkyl and benzyl; or R¹ and R² are taken together with the nitrogen to which they are attached and form

15



20

A² is selected from the group consisting of L-His, L-Cys, L-Cys(Me), L-Phe, L-Phe-R³, L-Val, L-Ala, L-Ile, L-Leu and L-Tyr;

A³ is selected from the group consisting of L-Val, L-Leu, L-Ile, L-Tyr, L-Phe and L-Phe-R³;

A⁴ is selected from the group consisting of a covalent bond, L-Phe, L-Phe-R³, L-Tyr, and L-Leu;

wherein R³ is attached to the aromatic ring of phenylalanine and for each occurrence is selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ alkoxy, benzyl, fluoro, trifluoromethyl and chloro; and


Q¹ is selected from the group consisting of t-butoxycarbonyl, benzyloxycarbonyl, R⁴CO and phenylcarbonyl, wherein R⁴ is hydrogen, C₁-C₆ alkyl or benzyl.

2. A compound or a pharmaceutically acceptable base salt thereof according to claim 1 wherein A¹ is -NR¹R² and A⁴ is a covalent bond or L-Tyr.

35

-20-


3. A compound or a pharmaceutically acceptable base salt thereof according to claim 2 wherein A^2 is L-Phe, L-Val, L-Ala, L-Ile or L-Leu.
- 5 4. A compound or a pharmaceutically acceptable base salt thereof according to claim 3 wherein A^2 is L-Ala and A^3 is L-Val.
5. A compound or a pharmaceutically acceptable base salt thereof according to claim 4 wherein Q^1 is t-butoxycarbonyl.
6. A compound or a pharmaceutically acceptable base salt thereof
10 according to claim 5 wherein R^1 and R^2 are taken together and form



15

7. A compound or a pharmaceutically acceptable base salt thereof according to claim 6 wherein A^4 is L-Tyr.
- 20 8. A compound or a pharmaceutically acceptable base salt thereof according to claim 6 wherein A^4 is a covalent bond.
9. A compound of the formula

25




- 30 wherein A^1 is L-Pro- NR^1R^2 or $\text{-NR}^1\text{R}^2$, where R^1 and R^2 are independently selected from the group consisting of hydrogen, $\text{C}_1\text{-C}_6$ alkyl and benzyl; or R^1 and R^2 are

35

-21-


taken together with the nitrogen to which they are attached and form




10
 A^2 is selected from the group consisting of L-His, L-Cys, L-Cys(Me), L-Phe, L-Phe- R^3 , L-Val, L-Ala, L-Ile, L-Leu and L-Tyr;

wherein R^3 is attached to the aromatic ring of the phenylalanine and is selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkoxy, benzyl, fluoro, trifluoromethyl and chloro.

15
 10. A compound of the formula



wherein A^1 is L-Pro- NR^1R^2 or NR^1R^2 , where R^1 and R^2 are independently selected from
 25 the group consisting of hydrogen, C_1 - C_6 alkyl and benzyl; or R^1 and R^2 are taken together with the nitrogen to which they are attached and form



A^2 is selected from the group consisting of L-His, L-Cys, L-Cys(Me), L-Phe, L-Phe- R^3 ,
 35 L-Val, L-Ala, L-Ile, L-Leu and L-Tyr;

-22-

A³ is selected from the group consisting of L-Val, L-Leu, L-Ile, L-Tyr, L-Phe and L-Phe-R³;


5 A⁴ is selected from the group consisting of a covalent bond, L-Phe, L-Phe-R³, L-Tyr, and L-Leu;

wherein R³ is attached to the aromatic ring of the phenylalanine and for each occurrence is selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ alkoxy, benzyl, fluoro, trifluoromethyl and chloro; and

10 Q¹ is selected from the group consisting of t-butoxycarbonyl, benzyloxycarbonyl, R⁴CO and phenylcarbonyl, wherein R⁴ is hydrogen, C₁-C₆ alkyl or benzyl.


11. A compound of the formula

15



wherein A¹ is L-Pro-NR¹R² or -NR¹R², where R¹ and R² are independently selected from
 20 the group consisting of hydrogen, C₁-C₆ alkyl and benzyl; or R¹ and R² are taken together with the nitrogen to which they are attached and form

25



A² is selected from the group consisting of L-His, L-Cys, L-Cys(Me), L-Phe, L-Phe-R³,
 30 L-Val, L-Ala, L-Ile, L-Leu and L-Tyr;

A³ is selected from the group consisting of L-Val, L-Leu, L-Ile, L-Tyr, L-Phe and L-Phe-R³;

A⁴ is selected from the group consisting of a covalent bond, L-Phe, L-Phe-R³, L-Tyr, and L-Leu;

35

-23-

wherein R^3 is attached to the aromatic ring of the phenylalanine and for each occurrence is selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkoxy,

5 benzyl, fluoro, trifluoromethyl and chloro; and

Q^1 is selected from the group consisting of t-butoxycarbonyl, benzyloxycarbonyl, R^4CO and phenylcarbonyl, wherein R^4 is hydrogen, C_1 - C_6 alkyl or benzyl.

12. A pharmaceutical composition comprising an amount of a compound or a pharmaceutically acceptable base salt thereof according to claim 1 and a
10 pharmaceutically acceptable diluent or carrier.

13. A method of inhibiting interleukin 1β converting enzyme (ICE) in a mammal in need thereof which comprises administering to said mammal an interleukin 1β converting enzyme inhibiting amount of a compound or a pharmaceutically acceptable base salt thereof according to claim 1.

15 14. A method of treating an antiinflammatory condition in a mammal which comprises administering to said mammal an antiinflammatory amount of a compound or a pharmaceutically acceptable base salt thereof according to claim 1.

20

25

30

35

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 93/03589

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C07K5/02;	C07K7/02;	C07D307/33; A61K37/64
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C07K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ^o	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
P,X	JOURNAL OF ORGANIC CHEMISTRY vol. 57, no. 26, 1992, EASTON US pages 7309 - 7314 R.P. TOBINSON AND K.M. DONAHUE 'Synthesis of a Peptidyl Difluoro Ketone Bearing the Aspartic Acid Side Chain: An Inhibitor of Interleukin-1beta Converting Enzyme' see the whole document ---	1-14
A	WO,A,9 115 577 (IMMUNEX CORPORATION) 17 October 1991 see page 3, line 25 - line 35 see page 4, line 31 - page 6, line 2; claim 14; example 29 --- -/--	1,12-14
^o Special categories of cited documents : ¹⁰ "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
19 AUGUST 1993	23 -09- 1993	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	FUHR C.K.	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	EP,A,0 195 212 (MERRELL DOW PHARMACEUTICALS INC.) 24 September 1986 see page 53, line 1 - page 56, line 5; claims -----	1,12-14

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/03589

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark : Although claims 13, 14 are directed to a method of treatment of the human/animal body the search has been carried out and based on the alleged effects of the compound.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest☐ The additional search fees were accompanied by the applicant's protest.☐ No protest accompanied the payment of additional search fees.

US 9303589
SA 73325

19/08/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9115577	17-10-91	AU-A- 7775991	30-10-91
EP-A-0195212	24-09-86	AU-B- 600226	09-08-90
		AU-A- 5288186	07-08-86
		JP-A- 61183253	15-08-86

THIS PAGE BLANK (USPTO)